

Update- ATSDR Policy Guideline for Dioxins and Dioxin-Like Compounds in Residential Soil

The Agency for Toxic Substances and Disease Registry (ATSDR) has updated its *Policy Guideline for Dioxins and Dioxin-Like Compounds in Residential Soil*. The most significant change is the deletion of the 1-part per billion (ppb) action level criteria. This update does *not* reflect a change in the hazard assessment that established ATSDR's 1998 Toxicological Profile and Minimum Risk Level (MRL) determination of the health concerns associated with dioxin exposure. The revised policy reflects ATSDR's concern that the previously established *action level* had been misinterpreted.

STATEMENT OF ISSUES

In 1998, the Agency for Toxic Substances and Disease Registry (ATSDR) adopted a Final Policy Guideline for Dioxin and Dioxin-Like Compounds (De Rosa et al. 1999a). The 1998 policy guideline was accompanied by a Technical Support Document for ATSDR Policy Guideline (De Rosa et al. 1999b). The initiative to develop this policy guideline was based on a request from the U.S. Environmental Protection Agency (EPA) to evaluate the protectiveness of the EPA Superfund Policy for Dioxins in Residential Soils, which established 1 part per billion (ppb) (1,000 parts per trillion [ppt]) total dioxin toxicity equivalents (TEQ) as the starting point for making clean-up decisions. In addition, the 1998 policy guideline was to provide guidance to health assessors in evaluating the public health implications of dioxin and dioxin-like compounds (e.g., 2,3,7,8-tetrachlorodibenzo-*p*-dioxin [TCDD], chlorinated dibenzodioxins [CDDs], chlorinated dibenzofurans [CDFs], and other structurally related groups of chemicals from the family of halogenated aromatic hydrocarbons) in residential soils near or on hazardous waste sites. As stated in the 1998 document, "these guidelines and procedures apply to human exposure for **direct** ingestion of soils contaminated with dioxin and dioxin-like compounds in residential areas and may not be appropriate for other exposure scenarios."

The 1998 Policy Guideline established a screening level (≤ 0.05 ppb TEQ), an evaluation level (> 0.05 ppb TEQ, < 1 ppb), and an action level (≥ 1 ppb TEQ) for dioxins in residential soil and made recommendations for specific considerations or public health actions.

Problems Regarding the 1998 Policy Guideline

Differences in the interpretation of the 1998 guideline have led to inconsistencies in the application of the screening and action levels to public health assessments. One problem has been the concept of an "action level" for public health activities. The action level has been interpreted by others as: 1) a soil concentration that defines a public health hazard, 2) an ATSDR clean-up level which implies the need for site remediation, or 3) an initial screening level that defines a safe level of exposure, below which there is no public health concern. None of these interpretations were either intended or appropriate.

In addition, the 1998 ATSDR action level triggered a set of recommendations for “potential public health actions”, including surveillance, research, health studies, community education, and exposure investigations. However, these actions should be activities to be considered by health assessors regardless of exceeding a 1 ppb level in soil. The 1998 ATSDR policy guideline established an *action level*, yet did not require that specific actions be conducted.

ATSDR has established environmental screening values for chemicals to be used by health assessors to assess exposures. No other chemical has an action level as was established for dioxins in soil in the 1998 policy guideline. This inconsistency alone has led to confusion regarding the appropriate screening value for soil dioxin levels.

OBJECTIVES OF REVISIONS TO THE DIOXIN SOIL POLICY GUIDELINE

ATSDR is updating the soil policy guideline for assessing exposure to dioxin and dioxin-like compounds in soil. Details about the health concerns for dioxins have not been revised as summarized in the Technical Support Document for ATSDR Final Policy Guideline: Dioxin and Dioxin-Like Compounds in Soil (De Rosa et al. 1999b) and in the ATSDR Toxicological Profile for Chlorinated Dibenzo-*p*-Dioxins (ATSDR 1998).

The 2005 ATSDR Dioxin and Dioxin-Like Compounds Soil Policy Guideline ...

- **Deletes the 1 ppb action level as the criteria for taking specific public health actions.**

The 1-ppb dioxin soil concentration should not be used as a comparison value for defining public health hazards in public health assessments and consultations. The 1-ppb action level can be cited by health assessors as the Superfund Dioxin Cleanup policy criteria (EPA 1989, 1998).

- **Retains the 0.05 ppb screening level.**

The minimal risk level (MRL)-based environmental media evaluation guide (EMEG) of 0.05 ppb for dioxin TEQ in soil is retained as the basis for screening soil concentrations. Levels exceeding this screening level should be evaluated as described in the *ATSDR Public Health Assessment Guidance Manual (PHAGM)* (ATSDR 2005). This clarification will ensure that evaluation of dioxins and dioxin-like compounds in soil will be done in the same manner as all other soil contaminants.

- **Recommends exposure pathways analyses for dioxins and dioxin-like compounds.**

The focus of the guideline is the assessment of direct exposure to soil contamination, **particularly** soil ingestion. However, health assessors should be aware of the potential impact of indirect exposure pathways on exposed populations in site-specific health assessments. This document does not provide specific guidance on how these indirect pathways should be assessed. However,

the PHAGM document does provide assistance in evaluating indirect exposure pathways such as food chain contamination (ATSDR 2005).

UPDATED GUIDANCE FOR ASSESSING DIOXINS IN RESIDENTIAL SOILS

This update does *not* reflect a change in the assessment of health hazards associated with dioxin exposure as summarized in the 1998 ATSDR Toxicological Profile and in the derivation of the Minimum Risk Level (MRL). The revised policy reflects ATSDR's concern that the previously established *action level* had been misinterpreted. ATSDR continues to review the scientific literature regarding dioxin toxicity to ensure that the MRL is protective of human health.

To identify chemicals that may present at levels of potential health concern at hazardous waste sites, ATSDR's PHAGM describes the process for evaluating environmental sampling data for soil contaminants, including identification of chemicals of concern by comparing their concentration to a screening value or EMEG, characterization of exposure pathways and magnitude of exposure, and evaluation of public health hazard.

Step 1: Comparison of Environmental Data to ATSDR Screening Value

Studies in animals have demonstrated a wide range of health effects associated with dioxin exposure including death, cancer, and wasting, as well as hepatic, immunologic, neurologic, endocrine, reproductive, and developmental effects (ATSDR 1998, De Rosa et al. 1999b). An extensive body of research suggests that dioxins and DLCs are likely to act through a common mechanism of action to induce toxicity (ATSDR 1998). Based on this understanding, the relative potency of a number of DLCs compared to TCDD has been used to derive Toxicity Equivalent Factors (TEFs) for specific DLCs, as developed by the World Health Organization (Van Den Berg 1998). ATSDR uses TEFs to summarize the dioxin content of a soil sample as the total dioxin toxicity equivalents (TEQs), as described in the appendices and in the ATSDR Toxicological Profile for Chlorinated Dibenzo-p-Dioxins (ATSDR 1998).

The chronic-duration oral MRL of 1 picogram/kilogram body weight/day (pg/kg/day) for TCDD, or total TEQ, (ATSDR 1998) is based on neurobehavioral effects in monkeys (Schantz et al. 1992). A MRL is an estimate of the daily human exposure to a hazardous substance that is likely to be without appreciable risk of adverse non-cancer health effects over a specified duration and route of exposure (ATSDR 1996). Uncertainty factors of 90 (total) were used in calculations of the MRL (for additional details, see ATSDR 1998). This value is consistent with the tolerable daily intake (TDI) of 1-4 pg/kg/day for TCDD or total TEQs (WHO 1998), the TDI of 4 pg TEQs/kg/day (Japanese Environmental Agency 1999), and the tolerable weekly intake of 7 pg/kg/week of TCDD of total TEQs (European Union 2000).

Using the chronic MRL value (1 pg/kg/day), exposure assumptions for a child (soil ingestion rate = 200 mg/day and 10 kg body weight), and assumption of 100% bioavailability, an EMEG of **0.05 ppb** (50 ppt) TEQ was derived as a screening value for residential soils contaminated with dioxins and dioxin-like compounds.

